


Showcasing research from Tatsuya Nabeshima's Laboratory/Tsukuba Research Center for Interdisciplinary Materials Science (TIMS), Graduate School of Pure and Applied Sciences, University of Tsukuba, Japan.

Photoisomerization locking of azobenzene by formation of self-assembled macrocycle

Self-assembled ring- and cage-shaped complexes of an azobenzene-linked biscatecholate ligand were synthesized, and these complexes were inert to photoisomerization. This fact indicates that the self-assembly approach is an efficient and new method for the photoisomerization locking of azobenzenes.

As featured in:



See Tatsuya Nabeshima *et al.*,
Chem. Commun., 2012, **48**, 5724.

RSC Publishing

www.rsc.org/chemcomm

Registered Charity Number 207890

Cite this: *Chem. Commun.*, 2012, **48**, 5724–5726

www.rsc.org/chemcomm

COMMUNICATION

Photoisomerization locking of azobenzene by formation of a self-assembled macrocycle†

Masaki Yamamura, Yuki Okazaki and Tatsuya Nabeshima*

Received 22nd December 2011, Accepted 27th January 2012

DOI: 10.1039/c2cc18014f

Reaction of an azobenzene-linked biscatechol ligand with boron and titanium sources gave ring- and cage-shaped complexes in a self-assembly fashion, respectively. These complexes were inert to photoisomerization though the ligand itself was isomerized upon photoirradiation. The self-assembled macrocyclization caused inhibition of the photoisomerization.

Photochromic compounds are fascinating materials because of their application in optical memory devices, molecular machines, and switchable molecular receptors.¹ On the other hand, locking of photoreaction has been investigated because the inhibition plays a critical role in the design of highly stable dyes,² photorefractive materials,³ and luminescent dyes.⁴ More recently, a combination of photo and other external stimuli was reported to lead to multi-responsive systems.^{5,6} In these multi-responsive systems, photoreaction controlled by an external effector, such as a proton,⁷ ion,⁸ Lewis acid,⁹ or electron,¹⁰ provides molecular switches serving as a dual-input logic system and nondestructive molecular memory based on locked and unlocked photoreactions by the addition of the effector. Here, we focused on a self-assembly approach for the photoisomerization locking by forming a rigid macrocyclic azobenzene (Fig. 1). Azobenzene, one of the most commonly

used photochromic units, features a light-induced *trans*–*cis* isomerization. Locking of the photoswitch on supramolecular macrocycles having photochromic units has potential for tuning the isomerization behaviour due to reversible formation of the macrocyclic structure. Isomerization of macrocyclic azobenzene derivatives inhibited by a guest or in a condensed state was reported;⁸ perfect locking of isomerization by self-assembled macrocyclization has not been reported as far as we know.

Herein, an azobenzene-linked biscatechol ligand was designed and synthesized for self-assembled rigid macrocycle formation. The macrocyclic complexes of azobenzene-biscatechol showed locking of the photoisomerization behaviour.

The azobenzene-linked biscatechol ligand *trans*-LH₄ (Scheme 1) was synthesized as follows: the cross-coupling reaction of 4,4'-dibromoazobenzene with 2,3-dimethoxyphenylboronic acid in the presence of Pd(PPh₃)₄ gave a precursor, 4,4'-bis(2,3-dimethoxyphenyl)azobenzene (72% yield), which was demethylated by treatment with BBr₃ to give *trans*-LH₄ (75% yield).† The structural analysis of *trans*-LH₄ by X-ray crystallography revealed the usual bond lengths and angles as seen in *trans*-azobenzene.¹¹ In the UV-vis spectrum, absorption maxima





Fig. 1 Control of photoisomerization behaviour of azobenzene by self-assembly approach.

Tsukuba Research Center for Interdisciplinary Materials Science (TIMS), Graduate School of Pure and Applied Sciences, University of Tsukuba, 1-1-1, Tennodai, Tsukuba, Ibaraki 305-8571, Japan. E-mail: nabeshima@chem.tsukuba.ac.jp; Fax: +81 29-853-4507; Tel: +81 29-853-4507

† Electronic supplementary information (ESI) available: The detailed synthetic procedure, X-ray crystallographic analysis, 1D NMR spectra, DOSY, ESI- and MALDI-TOF-MS, and computational study. CCDC 855272. For ESI and crystallographic data in CIF or other electronic format see DOI: 10.1039/c2cc18014f



Scheme 1 Self-assembly formation of azobenzene-linked ring [*L*₄B₄]⁴⁻ and cage [*L*₃Ti₂]⁴⁻.

of $\pi-\pi^*$ and $n-\pi^*$ transitions appeared at 360 and 440 nm, respectively.

Two different types of complexes of *trans*-LH₄ bearing boron and titanium catecholate, which are known as a joint part of ring-¹² and cage-shaped¹³ self-assemblies of biscatecholate ligands, respectively, were synthesized to construct the macrocyclic structure. First, the borate complex was synthesized by the reaction of *trans*-LH₄ with triisopropyl borate in the presence of Et₃N to give a red glassy solid in 88% isolated yield (Scheme 1). ¹H, ¹³C and ¹¹B NMR spectra in DMSO-*d*₆ showed highly symmetric signals assigned to one set of the biscatecholate boron and symmetric azobenzene moieties with Et₃NH⁺ as a counter cation (Fig. S2, ESI†). All the protons were characterized by ¹H-¹H COSY and NOESY. When (¹¹Bu₄N)OH was used instead of Et₃N, the product showed the same NMR signals in the aromatic region. No influence of the counter cation on NMR indicates the separation of an ion pair in solution. The elemental analysis corresponds to the cyclic oligomer [L_{*n*}B_{*n*}](Et₃NH)_{*n*}. In addition, negative mode ESI-TOF-MS measurement also confirmed the formation of the cyclic compound by observation of an ion peak corresponding to [L_{*n*}B_{*n*}]^{*n*-} at *m/z* of 405.1 as a main peak and no acyclic oligomer was observed. The isotopic pattern of the ion peak correlated with the theoretical distribution of tetramer [L₄B₄]⁴⁻ mixed with trimer [L₃B₃]³⁻ is shown in Fig. S7, ESI.† A diffusion-ordered NMR spectroscopic (DOSY) experiment strongly supported a single discrete product by the alignment of a single set of diffusion coefficient peaks for the product (Fig. S5, ESI†). The diffusion constant was determined to be $7.06 \times 10^{-11} \text{ m}^2 \text{ s}^{-1}$, and the corresponding hydrodynamic radius, *R*_h, was a much larger value, 15.6 Å, than that of *trans*-LH₄, 4.88 Å. A molecular modelling study was performed for the macrocyclic complexes by DFT calculation.¹⁴ There are four diastereomers in the macrocycle [(*trans*-L)₄B₄]⁴⁻ due to the chirality of the boron catecholate moieties (Fig. S15, ESI†). The modelling study of the four diastereomers suggests that the most stable isomer is the *D*₄ symmetric one with its four boron centres having the same chirality. The observed *R*_h was comparable to the molecular size of the calculated model structure of tetramer [(*trans*-L)₄B₄]⁴⁻ (15.5 Å) rather than trimer [L₃B₃]³⁻ (12.6 Å, Fig. S14, ESI†). Trimer [L₃B₃]³⁻ observed in ESI-MS should exist as only a minor species with a very small population to be observed in the NMR spectra. Thus, the ring-shaped self-assembled complex [(*trans*-L)₄B₄]⁴⁻ was efficiently formed.

The titanium complex was quantitatively synthesized by the reaction of *trans*-LH₄ with Ti(O)(acac)₂ in the presence of Na₂CO₃ (99% yield, Scheme 1). The ¹H NMR spectrum showed only one discrete product (Fig. S3, ESI†). The MALDI-TOF-MS, DOSY experiment, and the model study by DFT calculation suggested the formation of the *D*₃ symmetric trigonal-pyramidal cage-shaped complex [(*trans*-L)₃Ti₂]⁴⁻ (Fig. S16, ESI†),¹⁵ similarly to reported cage-shaped triscatecholate Ti complexes.¹³ To prove the detailed macrocyclic effect, acyclic boron and titanium complexes [L'₂B](Et₃NH) and [L'₃Ti]Na₂ were synthesized following a procedure similar to those for [(*trans*-L)₄B₄]⁴⁻ and [(*trans*-L)₃Ti₂]⁴⁻,¹⁶ respectively (Chart 1). Finally, synthesis of the self-assembled complexes of *cis*-LH₄ was attempted toward photoisomerization locking of the *cis*-azobenzene. But the reaction of *cis*-LH₄ with boron and titanium sources only gave a mixture of unidentified compounds




Chart 1 Acyclic complexes [L'₂B]⁻ and [L'₃Ti]²⁻.

and the *trans*-isomers, [(*trans*-L)₄B₄]⁴⁻ and [(*trans*-L)₃Ti₂]⁴⁻, respectively.

The photoisomerization study of *trans*-LH₄, [(*trans*-L)₄B₄]⁴⁻ and [(*trans*-L)₃Ti₂]⁴⁻ was performed using a 400 W high-pressure mercury lamp. Irradiation of UV light (360 nm) onto a DMSO solution of *trans*-LH₄ resulted in a decrease in λ_{max} at 360 nm and an increase at 500 nm in the UV-vis spectra with isosbestic points. This spectral change corresponds to the typical *trans/cis* isomerization of azobenzene. The ¹H NMR spectrum showed new peaks assigned to the *cis*-isomer after irradiation, and the *trans/cis* ratio is 56/44 in the photostationary state at 360 nm. Irradiation of visible light (440 nm) onto the *cis/trans* mixture of LH₄ resulted in recovery of the absorption of the *trans*-form, which represents reversible isomerization. In the UV-vis spectra of [(*trans*-L)₄B₄]⁴⁻ in DMSO, a broad absorption band was observed around 400 nm, which represents a $\pi-\pi^*$ transition overlapped with an $n-\pi^*$ band. The λ_{max} was red-shifted over 30 nm compared to *trans*-LH₄. This shift is due to the electron-donation of the catecholate complex. The isomerization experiment of [(*trans*-L)₄B₄]⁴⁻ was carried out by irradiation at different wavelengths (360 nm and 440 nm). However, no spectral change was observed even after prolonged photoirradiation for 60 min (Fig. 2). The inhibition of photoisomerization is based on either the conformational rigidity as shown in azobenzene-linked rigid cyclophanes¹⁷ or the substitution effect of the catecholate complex unit. The photoisomerization of the *trans*-form of acyclic complex [L'₂B]⁻ was evaluated under the same conditions, and isomerization was confirmed by UV-vis and NMR spectra; the reaction reached the photostationary state after irradiation at 360 nm for 60 s (Fig. 2). The ratio of isomers in the photostationary state was determined to be 35 : 65 (*trans/cis*)




Fig. 2 UV-vis spectral change of [L'₂B](Et₃NH) (a dotted line, right axis) and [L₄B₄](Et₃NH)₄ (a solid line, left axis) after photoirradiation with a high-pressure mercury lamp (360 nm) in DMSO.




Fig. 3 UV-vis spectral change of $[\text{L}'_3\text{Ti}]\text{Na}_2$ (a dotted line) and $[\text{L}_3\text{Ti}_2]\text{Na}_4$ (a solid line) after photoirradiation with a high-pressure mercury lamp (360 nm) in DMSO.

by the integral ratio in the NMR spectrum. This result is in contrast with the inert nature of macrocycle $[(\text{trans-L})_4\text{B}_4]^{4-}$ to photoirradiation. The locking of photoisomerization of $[(\text{trans-L})_4\text{B}_4]^{4-}$ is ascribed to the molecular rigidity of the macrocyclic structure.

In the UV-vis spectra, the broad shoulder peaks of $[(\text{trans-L})_3\text{Ti}_2]^{4-}$ and $[(\text{trans-L}')_3\text{Ti}]^{4-}$, overlapped with the absorptions of azobenzene, reached the region of over 600 nm assigned to the MLCT bands (Fig. 3). Cage-shaped titanium complex $[(\text{trans-L})_3\text{Ti}_2]^{4-}$ did not isomerize upon photoirradiation. In contrast, photoirradiation of acyclic $[(\text{trans-L}')_3\text{Ti}]^{4-}$ resulted in a small but detectable spectral change. The spectrum was recovered by allowing the sample to stand in the dark, indicating the thermal back-isomerization of the *cis*-isomer of $[\text{L}'_3\text{Ti}]^{2-}$. A prolonged photoirradiation time or heating of $[(\text{trans-L}')_3\text{Ti}]^{2-}$ produced decomposed products. Conversely, cage-shaped $[(\text{trans-L})_3\text{Ti}_2]^{4-}$ did not change under the same conditions, revealing the photoisomerization locking and high stability of $[(\text{trans-L})_3\text{Ti}_2]^{4-}$. In addition, the NMR study also showed no isomerization of $[(\text{trans-L})_3\text{Ti}_2]^{4-}$ and isomerization of $[(\text{trans-L}')_3\text{Ti}]^{4-}$. Inhibition of the isomerization of $[(\text{trans-L})_3\text{Ti}_2]^{4-}$ should be ascribed to the macrocyclic structure.

In summary, the azobenzene-linked ring $[(\text{trans-L})_4\text{B}_4]^{4-}$ and cage $[(\text{trans-L})_3\text{Ti}_2]^{4-}$ were efficiently synthesized by a coordination-driven self-assembly of the azobenzene-linked biscatechol ligand *trans-LH*₄. The acyclic azobenzenes bearing catecholate moieties *trans-LH*₄, $[(\text{trans-L}')_2\text{B}]^-$, and $[(\text{trans-L}')_3\text{Ti}]^{2-}$ were isomerized by photoirradiation though the photoisomerization of the self-assembled macrocyclic azobenzenes $[(\text{trans-L})_4\text{B}_4]^{4-}$ and $[(\text{trans-L})_3\text{Ti}_2]^{4-}$ did not proceed at all. This fact indicates that the self-assembly approach is an efficient and new method for the photoisomerization locking of azobenzenes. Now we are investigating the application of this strategy for construction of more sophisticated molecular switches.

This research was financially supported by Grants-in-Aid for Scientific Research from the Ministry of Education, Culture, Sports, Science, and Technology of Japan.

Notes and references

- 1 *Molecular Switches*, ed. B. L. Feringa, Wiley-VCH, Weinheim, 2001; D. Bléger, Z. Yu and S. Hecht, *Chem. Commun.*, 2011, **47**, 12260–12266.
- 2 J. Griffiths, *Chem. Soc. Rev.*, 1972, **1**, 481–493.
- 3 O. Ostroverkhova and W. E. Moerner, *Chem. Rev.*, 2004, **104**, 3267–3314.
- 4 M. Shimomura and T. Kunitake, *J. Am. Chem. Soc.*, 1987, **109**, 5175–5183; M. Han and M. Hara, *J. Am. Chem. Soc.*, 2005, **127**, 10951–10955; J. Yoshino, N. Kano and T. Kawashima, *Chem. Commun.*, 2007, 559–561.
- 5 H. Nishihara, *Coord. Chem. Rev.*, 2005, **249**, 1468–1475; M. Akita, *Organometallics*, 2011, **30**, 43–51.
- 6 S.-S. Sun and A. J. Lees, *J. Am. Chem. Soc.*, 2000, **122**, 8956–8967; S.-S. Sun, J. A. Anspach and A. J. Lees, *Inorg. Chem.*, 2002, **41**, 1862–1869; C. S. Pecinovsky, E. S. Hatakeyama and D. L. Gin, *Adv. Mater.*, 2008, **20**, 174–178.
- 7 M. Irie, O. Miyatake and K. Uchida, *J. Am. Chem. Soc.*, 1992, **114**, 8715–8716; H. M. D. Bandara, T. R. Friss, M. M. Enriquez, W. Isley, C. Incarvito, H. A. Frank, J. Gascon and S. C. Burdette, *J. Org. Chem.*, 2010, **75**, 4817–4827.
- 8 M. Takeshita, C. F. Soong and M. Irie, *Tetrahedron Lett.*, 1998, **39**, 7717–7720; H.-S. Tang, N. Zhu and V. W.-W. Yam, *Organometallics*, 2007, **26**, 22–25; X. Li, Y. Ma, B. Wang and G. Li, *Org. Lett.*, 2008, **10**, 3639–3642; S. Venkataramani, U. Jana, M. Dommaschk, F. D. Sönnichsen, F. Tuzek and R. Herges, *Science*, 2011, **331**, 445–448.
- 9 Y. Wu, S. Chen, Y. Yang, Q. Zhang, Y. Xie, H. Tian and W. Zhu, *Chem. Commun.*, 2012, **48**, 528–530.
- 10 Y. F. Liu, C. Lagrost, K. Costuas, N. Touchar, H. Le Bozec and S. Rigaut, *Chem. Commun.*, 2008, 6117–6119; K. Motoyama, T. Koike and M. Akita, *Chem. Commun.*, 2008, 5812–5814; Y. Tanaka, T. Ishisaka, A. Inagaki, T. Koike, C. Lapinte and M. Akita, *Chem.–Eur. J.*, 2010, **16**, 4762–4776.
- 11 Crystallographic data for $\text{LH}_4\cdot\text{thf}_2$: $\text{C}_{32}\text{H}_{34}\text{N}_2\text{O}_6$, $M_w = 542.61$, triclinic, $P\bar{1}$, $a = 8.9085(12)$, $b = 11.6096(15)$, $c = 14.9458(19)$ Å, $\alpha = 69.838(2)^\circ$, $\beta = 74.1440(10)^\circ$, $\gamma = 85.913(2)^\circ$, $V = 1395.4(3)$ Å³, $Z = 2$, $D_{\text{calcd}} = 1.291$ g cm⁻³, $\lambda = 0.71073$ Å, $\mu = 0.089$ mm⁻¹, $T = 120(2)$ K, 6662 reflections, 4796 unique ($R_{\text{int}} = 0.0109$), $\text{GOF} = 1.015$, $R_1 = 0.0360$ ($I > 2\sigma(I)$), $wR_2 = 0.1064$ (all data), CCDC 855272.
- 12 B. F. Abrahams, D. J. Price and R. Robson, *Angew. Chem., Int. Ed. Engl.*, 2006, **45**, 806–810; B. F. Abrahams, B. A. Boughton, H. Choy, O. Clarke, M. J. Grannas, D. J. Price and R. Robson, *Inorg. Chem.*, 2008, **47**, 9797–9803; H. Danjo, K. Hirata, S. Yoshigai, I. Azumaya and K. Yamaguchi, *J. Am. Chem. Soc.*, 2009, **131**, 1638–1639.
- 13 M. Albrecht and S. Kotila, *Angew. Chem., Int. Ed. Engl.*, 1995, **34**, 2134–2137; M. Albrecht, H. Rottele and P. Burger, *Chem.–Eur. J.*, 1996, **2**, 1264–1268; D. L. Caulder, C. Brückner, R. E. Powers, S. König, T. N. Parac, J. A. Leary and K. N. Raymond, *J. Am. Chem. Soc.*, 2001, **123**, 8923–8938; M. Albrecht, I. Jansera and R. Fröhlich, *Chem. Commun.*, 2005, 157–165.
- 14 Firstly, equilibrium conformations were determined by Monte-Carlo method at the PM3 level. Then, each equilibrium conformer was optimized at the BLYP/6-31G* level.
- 15 In the negative mode MALDI-TOF-MS of the product, a main ion peak is observed as an isotopic cluster centred at m/z 1303.1 that matches with the theoretical isotope distribution for $[\text{L}_3\text{Ti}_2 + \text{Na} + 2\text{H}]^-$. R_h of L_3Ti_2 in DMSO-*d*₆ was estimated to be 8.8 Å by DOSY experiment. The R_h value of $[\text{L}_3\text{Ti}_2]^{4-}$ is smaller than that of $[\text{L}_4\text{B}_4]^{4-}$ and well comparable with the calculated model structure.
- 16 $[\text{L}'_3\text{Ti}]^{2-}$ was obtained as a mixture of the *fac* and *mer* isomers because of their fast interconversion. See references for the interconversion of titanium triscatecholate complexes: A. Rosenheim, B. Raibmann and G. Z. Schendel, *Z. Anorg. Allg. Chem.*, 1931, **196**, 160–176; A. V. Davis, T. K. Firman, B. P. Hay and K. N. Raymond, *J. Am. Chem. Soc.*, 2006, **128**, 9484–9496.
- 17 R. Reuter and H. A. Wegner, *Chem. Commun.*, 2011, **47**, 12267–12276.